

SETS WITHOUT k -TERM PROGRESSIONS CAN HAVE MANY SMALLER PROGRESSIONS

JACOB FOX AND COSMIN POHOATA

ABSTRACT. Let $f_{s,k}(n)$ be the maximum possible number of s -term arithmetic progressions in a sequence $a_1 < a_2 < \dots < a_n$ of n integers which contains no k -term arithmetic progression. For all integers $k > s \geq 3$, we prove that

$$\lim_{n \rightarrow \infty} \frac{\log f_{s,k}(n)}{\log n} = 2,$$

which answers an old question of Erdős. In fact, we prove upper and lower bounds for $f_{s,k}(n)$ which show that its growth is closely related to the bounds in Szemerédi's theorem.

1. INTRODUCTION

Let $k \geq 3$ be an integer. In this paper, a k -term arithmetic progression of integers will denote as usual a set of the form $\{x, x + d, \dots, x + (k - 1)d\}$. If $d \neq 0$, then we say that the progression is *non-trivial*. If a set A does not contain any non-trivial k -term arithmetic progressions, we say that A is *k -AP free*. The study of k -AP free sets in the integers and other groups has been a central topic in additive combinatorics. Following the standard notation, we will denote by $r_k(n)$ the size of the largest k -AP free subset of $\{1, \dots, n\}$. The seminal result on this topic is Szemerédi's Theorem [10], which states that sets of integers with positive density contain arbitrarily long arithmetic progressions, or using the notation above $r_k(n) = o(n)$.

Since Szemerédi, the problem of finding better quantitative bounds for $r_k(n)$ has received a lot of attention, with impressive progress that led to many important tools, which in the meantime have become standard. For our application, we won't need the best bounds for each k , so we will limit ourselves to only mentioning Gowers' theorem [4, 5] that for each $k \geq 3$ there exists an absolute constant $c_k > 0$ such that

$$r_k(n) \ll \frac{n}{(\log \log n)^{c_k}}. \quad (1.1)$$

Regarding lower bounds, Rankin [8] showed that there exists a constant $c'_k > 0$ such that

$$r_k(n) \gg \frac{n}{2^{c'_k (\log n)^{1/\lceil \log k \rceil}}}. \quad (1.2)$$

Throughout the paper, all logarithms are base 2 and the signs \ll and \gg are the usual Vinogradov symbols.

Let $\mathcal{A}_k(n)$ be the set of n -term nonnegative integer sequences which contain no k -term arithmetic progression as a subsequence. Furthermore, let $f_s(A)$ denote the number of s -term arithmetic progressions in A , and finally let $f_{s,k}(n) = \max_{A \in \mathcal{A}_k(n)} f_s(A)$. In [3, page 119], Erdős observed that

$$\frac{\log f_{3,4}(n)}{\log n} > 1.4649$$

holds for infinitely many n by constructing examples of sequences $A \in \mathcal{A}_4(3^s)$ for which $f(A) = 3^{s-1}$. Furthermore, he noticed that for each $k > 3$ the limit $\lim_{n \rightarrow \infty} \log f_{3,k}(n) / \log n := f_{3,k}$ exists, and asked whether or not $f_{3,k}$ is always less than 2. In [1], Simmons and Abbott improved on Erdős' observation by showing that $f_{3,4}(n) \geq n^{1.623}$ holds infinitely often, and also proved that $s_{3,k} \rightarrow 2$ as k goes to infinity. Nonetheless, in the regime when k is fixed, there has been no further progress on understanding the limit $f_{3,k}$ as far as we are aware of. In this note, we settle Erdős' question in the negative by proving the following more general result.

Theorem 1.1. *For all integers $k > s \geq 3$, we have*

$$\lim_{n \rightarrow \infty} \frac{\log f_{s,k}(n)}{\log n} = 2.$$

In fact, we prove upper and lower bounds for $f_{s,k}(n)$ which show that its growth is closely related to the bounds in Szemerédi's theorem.

Theorem 1.2. *There exist absolute positive constants c and C such that, for integers $k > s \geq 3$ and every sufficiently large integer n , we have*

$$\left(\frac{c \cdot r_k(n)}{n} \right)^{2(s-2)} \cdot n^2 \leq f_{s,k}(n) \leq \left(\frac{r_k(n)}{n} \right)^C \cdot n^2.$$

In light of the bounds on $r_k(n)/n$ provided by (1.1) and (1.2), it is easy to check that Theorem 1.1 follows from Theorem 1.2; therefore, it suffices to prove the latter. We will do this already in Section 2. The proof of Theorem 1.2 will require a few ingredients from additive combinatorics, but we will state them in full as we will get to apply them, as they do not require much preparation.

Funding and acknowledgments. The first author was supported by a Packard Fellowship and by NSF grant DMS-1855635. The second author would like to thank Oliver Roche-Newton for helpful conversations.

2. PROOF OF THEOREM 1.2

We first prove the desired upper bound on $f_{s,k}(n)$. For $s \geq 3$, we have $f_{s,k}(n) \leq f_{3,k}(n)$, so in order to prove the upper bound it suffices to show that

$$f_{3,k}(n) \leq \left(\frac{r_k(n)}{n} \right)^C n^2$$

holds for some absolute constant $C > 0$ and sufficiently large n . We will in fact show this claim for $C = 1/25$. Let $A \in \mathcal{A}_k(n)$ and let pn^2 denote the number of three-term arithmetic progressions in A , where p is some positive real number (which is strictly less than 1); i.e. $f_3(A) = pn^2$.

To upper bound p , we will require the following variant of the Balog-Szemerédi-Gowers theorem (see [4] or [2]).

Theorem 2.1. *If A and B are sets of n integers and G is a bipartite graph between A and B with pn^2 edges such that partial sumset $A +_G B$ has size at most $K|A|$, then there is a subset A' of A with $|A'| \geq pn/4$ and*

$$|A' - A'| \ll K^4 p^{-5} n.$$

Here $A +_G B$ denotes as usual the sumset restricted to the edges coming from G , namely

$$A +_G B = \{a + b : a \in A, b \in B, (a, b) \in E(G)\}.$$

It is perhaps important to mention that Theorem 2.1 is a somewhat nonstandard version of the Balog-Szemerédi-Gowers theorem, which outputs directly a large set $A' \subset A$ with small difference set, without applying any Ruzsa-type inequality. We refer the reader to the proof of [2, Lemma 5.2, page 9], from which the following statement can also be extracted.

Lemma 2.2. *If a bipartite graph $G = (A, B, E)$ with $|A| = |B| = n$ has pn^2 edges, then there is a subset A' of A of size at least $pn/4$ such that every pair of vertices in A have at least $\Omega(p^5 n^3)$ paths of length four connecting them.*

Using Lemma 2.2, one can then deduce Theorem 2.1 in the usual way. Applied to the graph from the setup of Theorem 2.1, Lemma 2.2 produces $A' \subset A$ of size at least $pn/4$ such that every pair of vertices in A have at least $\Omega(p^5 n^3)$ paths of length four connecting them. This set happens to also satisfy $|A' - A'| \ll K^4 p^{-5} n$. Indeed, for each $a, a' \in A'$, consider a path of length four in G between them, say (a, b, a'', b', a') . For $y := a - a' \in A' - A'$, we can then write

$$a - a' = (a + b) - (a'' + b) + (a'' + b') - (a' + b') = x_1 - x_2 + x_3 - x_4,$$

where $x_1 = a + b$, $x_2 = a'' + b$, $x_3 = a'' + b'$, and $x_4 = a' + b'$ are all elements of $A +_G B$. Since for every $a, a' \in A'$ there are at least $\Omega(p^5 n^3)$ paths of length four between a and a' , this means every $y \in A' - A'$ can be written as $x_1 - x_2 + x_3 - x_4$ for at least $\Omega(p^5 n^3)$ quadruples $(x_1, x_2, x_3, x_4) \in (A +_G B)^4$. However, $|A +_G B| \leq Kn$ holds by assumption, so there are at most $K^4 n^4$ such quadruples. By the pigeonhole principle, it then follows that the number of distinct elements $y \in A' - A'$ is at most $O(K^4 p^{-5} n)$, as claimed.

Returning to the task of deriving the upper bound from Theorem 1.2, we apply Theorem 2.1 to the graph G where A and B are chosen to be two copies of our k -AP free A and with an edge between $(a, b) \in A \times A$ if $a + b = 2c$ for some $c \in A$. This graph has precisely pn^2 edges and we can apply Theorem 2.1 to it with $K = 1$ since

$$|A +_G A| = |\{2a : a \in A\}| = |A|.$$

This yields a subset $A' \subset A$ with $|A'| \geq p|A|/4$ and $|A' - A'| \ll p^{-5}n \ll p^{-6}|A'|$. At this point, we recall a version of the so-called Freiman-Ruzsa modelling lemma (see for instance [9, Theorem 2.3.5, page 127]).

Lemma 2.3. *Let S be a finite set of integers and let $r \geq 2$ be an arbitrary integer. Then, there is a set $S^* \subset S$ with $|S^*| \geq |S|/r^2$ which is Freiman r -isomorphic to a set of integers T such that*

$$T \subset \left\{ 1, 2, \dots, \left\lceil \frac{1}{r} \cdot |rS - rS| \right\rceil \right\}.$$

Here $rS - rS$ denotes the sumset $S + \dots + S - S - \dots - S$, where S appears $2r$ times. For the reader's convenience, we also recall that for any two commutative groups G_1, G_2 two sets $S \subset G_1$ and $T \subset G_2$ are said to be Freiman r -isomorphic if there exists a one to one map $\phi : S \rightarrow T$ such that for every $x_1, \dots, x_r, y_1, \dots, y_r$ in S (not necessarily distinct) the equation

$$x_1 + \dots + x_r = y_1 + \dots + y_r$$

holds if and only if

$$\phi(x_1) + \dots + \phi(x_r) = \phi(y_1) + \dots + \phi(y_r).$$

We combine Lemma 2.3 with (a consequence of) the classical Plünnecke-Ruzsa inequality, for which a simple proof can be found in [7].

Lemma 2.4. *Let S and T be finite sets of reals such that $|S + T| \leq \alpha|S|$, and let r, r' be positive integers. Then*

$$|rT - r'T| \leq \alpha^{r+r'}|S|.$$

Indeed, if we apply this with $S = A', T = -A', r = r' = 2$, and $\alpha = p^{-6}$, we have

$$|2A' - 2A'| \leq p^{-24}|A'| \leq p^{-24}n.$$

Therefore, by Lemma 2.3, there is a subset $A^* \subset A'$ with $|A^*| \gg pn$ which is Freiman 2-isomorphic to a set of integers $\phi(A^*)$ contained in the interval $\{1, \dots, \lceil p^{-24}n \rceil\}$. In particular, since ϕ preserves k -term arithmetic progressions,

$$pn \ll |A^*| = |\phi(A^*)| \leq r_k(\lceil p^{-24}n \rceil).$$

Lastly, recall that $r_k(n)$ is subadditive as a function of n , namely the inequality $r_k(n + n') \leq r_k(n) + r_k(n')$ holds for all positive integers n, n' . In particular, $r_k(\lceil p^{-24}n \rceil) \ll p^{-24}r_k(n)$, hence $pn \ll p^{-24}r_k(n)$, or equivalently $p^{25} \ll r_k(n)/n$. This means that A contains at most $(r_k(n)/n)^{1/25} n^2$ three-term arithmetic progressions. This completes the proof of the upper bound.

We next prove the desired lower bound on $f_{s,k}(n)$ in Theorem 1.2. We begin by revisiting some further simple properties of $r_k(n)$ as a function of n . In addition to being subadditive, we also recall that $r_k(n)$ is an increasing function, so $r_k(m) \leq r_k(n)$ if $m \leq n$. Together these imply that if $n \geq m$, we have $r_k(n) \leq \lceil \frac{n}{m} \rceil r_k(m) \leq \frac{2n}{m} r_k(m)$, so

$$\frac{r_k(n)}{2n} \leq \frac{r_k(m)}{m}. \quad (2.1)$$

For all positive integers m and n , we have

$$r_k(2mn) \geq r_k(m)r_k(n). \quad (2.2)$$

Indeed, if U is a subset of $\{1, \dots, m\}$ without a k -term arithmetic progression and V is a subset of $\{1, \dots, n\}$ without a k -term arithmetic progression, then the set

$$W = \{2u(n-1) + v : u \in U, v \in V\}$$

is a k -AP free subset of $\{1, \dots, 2mn\}$ of size $|U||V|$, so (2.2) follows.

In particular, if $n \geq N^{1/2}$, letting $m = \lfloor \frac{N}{2n} \rfloor$, we have

$$r_k(N) \geq r_k(2mn) \geq r_k(n)r_k(m) \geq r_k(n) \frac{m}{2n} r_k(n) \geq \frac{N}{8} \left(\frac{r_k(n)}{n} \right)^2,$$

where the first inequality follows from $r_k(n)$ being an increasing function, the second inequality is by (2.2), the third inequality is by (2.1) using $n \geq m$, and finally the fourth inequality is by substituting in $n \leq 4mN$. It thus follows that

$$\frac{r_k(N)}{N} \geq \frac{1}{8} \left(\frac{r_k(n)}{n} \right)^2. \quad (2.3)$$

Let $N = N_{n,k,s}$ be the least positive integer such that $r_k(N) = \lfloor n/s \rfloor$. Such an N exists since, for every m , $r_k(m+1) = r_k(m)$ or $r_k(m) + 1$ and $\lim_{m \rightarrow \infty} r_k(m) = \infty$. We will show that for $k > s \geq 3$ and n sufficiently large in terms of k , we have

$$f_{s,k}(n) \geq \left(\frac{n}{300sN} \right)^{s-2} n^2. \quad (2.4)$$

For n sufficiently large in terms of k , we have $n \geq N^{1/2}$ holds (for instance by (1.2)), so (2.3) implies that $n/N \geq s \cdot r_k(N)/N \geq s \cdot (1/8) \cdot (r_k(n)/n)^2$, and hence the lower bound from Theorem 1.2 follows from (2.4). We next prove (2.4) using a probabilistic construction of a k -AP free set A of n integers with many s -term arithmetic progressions.

For each $1 \leq i \leq s$, let d_i be an integer chosen uniformly and independently at random from the set $\{1, \dots, 2N\}$. Let $S \subset \{1, \dots, N\}$ be a k -AP free set of cardinality $r_k(N) = \lfloor n/s \rfloor$, and S_i denote the translate $\{x + 6(i-1)N - 1 + d_i : x \in S\}$, i.e. $S_i := S + \{6(i-1)N - 1 + d_i\}$.

Finally, let us consider the set $A \subset \{1, \dots, 6sN\}$ defined by

$$A := \bigcup_{i=1}^s S_i.$$

We first check that such a (random) set must be k -AP free. Indeed, the sets S_1, \dots, S_s are pairwise disjoint since, for each $1 \leq i \leq s$, we have

$$S_i \subset \{6(i-1)N + 1, \dots, 6(i-1)N + 3N - 1\}.$$

Furthermore, these sets are spaced out so that if an arithmetic progression contains an element from S_i and an element of S_j with $i \neq j$, then its common difference is at least $3N + 2$, in which case the arithmetic progression cannot

contain two elements in the same S_i . In particular, every arithmetic progressions in A of length longer than s must be a subset one of the S_i , and hence A is k -AP free. Finally, $|A| = s|S| = s\lfloor \frac{n}{s} \rfloor \leq n$, so A is indeed in $\mathcal{A}_k(n)$, or it can be artificially augmented to a set in $\mathcal{A}_k(n)$ by adding some elements that do not create k -term arithmetic progressions.

We next lower bound the expected number of s -term arithmetic progressions in A . The number of s -term arithmetic progressions $a, a + D, \dots, a + (s - 1)D$ with $a + (i - 1)D \in \{6(i - 1)N + N + 1, \dots, 6(i - 1)N + 2N\}$ for $1 \leq i \leq s$ is the same as the number of s -term arithmetic progressions in $\{1, \dots, N\}$ with any integer common difference, which is

$$N + 2 \sum_{a=1}^{N-1} \left\lfloor \frac{N-a}{s} \right\rfloor \geq \frac{1}{s} \binom{N}{2}.$$

For each such s -term arithmetic progression $a, a + D, \dots, a + (s - 1)D$ and for each sequence (a_1, \dots, a_s) of s elements from S , there is a choice of $d_1, \dots, d_s \in \{1, \dots, 2N\}$ such that $a_i + 6(i - 1)N - 1 + d_i = a + (i - 1)D$ for $1 \leq i \leq s$. Hence, the expected number of s -term arithmetic progressions in A is at least

$$\frac{1}{s} \binom{N}{2} |S|^s (2N)^{-s} \geq \frac{1}{4s} N^2 \left(\frac{\lfloor n/s \rfloor}{2N} \right)^s \geq \left(\frac{n}{300sN} \right)^{s-2} n^2.$$

Thus, there must exist a choice of such an A for which the number of s -term arithmetic progressions is at least this lower bound on the expected number, which completes the proof of (2.4) and hence Theorem 1.2.

REFERENCES

- [1] G. J. Simmons, H. L. Abbott, ‘How many 3-term arithmetic progressions can there be if there are no long ones?’, *The Amer. Math. Monthly*, Vol. 84, No. 8 (Oct., 1977), pp. 633-635.
- [2] J. Fox, B. Sudakov, ‘Dependent random choice’, *Random Structures Algorithms*, 38:6899, 2011.
- [3] P. Erdős, ‘Problems and results on combinatorial number theory’, Chapter 12 in *A survey of Combinatorial Theory* edited by J. Srivastava, North Holland, 1973.
- [4] W. T. Gowers, ‘A new proof of Szemerédi theorem for progressions of length four’, *Geom. Func. Anal.*, (8) 1998, 529-551.
- [5] W. T. Gowers, ‘A new proof of Szemerédi theorem’, *Geom. Func. Anal.*, (11) 2001, 465-588.
- [6] B. Green, T. Tao, ‘New bounds for Szemerédi’s theorem, III: A polylogarithmic bound for $r_4(N)$ ’, *Mathematika*, (63), Issue 3, 2017, 944-1040.
- [7] G. Petridis, ‘New proofs of Plünnecke-type estimates for product sets in groups’, *Combinatorica* 32 (2012), no. 6, 7.
- [8] R. A. Rankin, ‘Sets of integers containing not more than a given number of terms in arithmetical progression’, *Proc. Roy. Soc. Edinburgh*, Sect. A, (65) 1960/1961, 332-344.
- [9] I. Z. Ruzsa, ‘Sumsets and structure’, pp. 87-210 in *Combinatorial number theory and additive group theory*, edited by A. Geroldinger and I. Z. Ruzsa, Birkhäuser, Basel, 2009.
- [10] E. Szemerédi, ‘On sets of integers containing no k elements in arithmetic progression’, *Acta Arith.* 27 (1975), 199-245.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA 94305,
USA

E-mail address: `jacobfox@stanford.edu`

DEPARTMENT OF MATHEMATICS, CALIFORNIA INSTITUTE OF TECHNOLOGY, PASADENA,
CA 91106, USA

E-mail address: `apohoata@caltech.edu`